

Eggplant White Star

(*Solanum melongena*)

Germination

- Time of radicle emergence (3-6 days)
- Soil temperature 70-75°F (21-24°C).
- Keep media evenly moist but not saturated.
- Cover the seed lightly with coarse vermiculite.
- Soil pH 5.5-5.8 and soluble salts (EC) less than 0.5 mmhos/cm (2:1 extraction).
- Eggplant is very sensitive to high salts, particularly high ammonium, during germination.
- Keep ammonium levels to less than 10 ppm.

Plug Production

STAGE 1 - Time of radicle emergence (3-6 days)

- Soil temperature 70-75°F (21-24°C).
- Keep media evenly moist but not saturated.
- Cover the seed lightly with coarse vermiculite.
- Soil pH 5.5-5.8 and soluble salts (EC) less than 0.5 mmhos/cm (2:1 extraction).
- Eggplant is very sensitive to high salts, particularly high ammonium, during germination.
- Keep ammonium levels to less than 10 ppm.

STAGE 2 - Stem and cotyledon emergence (7-10 days)

- Soil temperature 70-75°F (21-24°C).
- Reduce moisture levels once radicle emergence occurs! Allow the soil to dry out slightly before watering for best germination and rooting.
- Keep soil pH 5.5-5.8 and EC to less than 0.5 mmhos/cm.
- Keep ammonium levels to less than 10 ppm.
- Begin fertilizing with 50 75 ppm N from 14 0 14 or 15-5-15 or a calcium/potassium nitrate feed once cotyledons are fully expanded.
- Alternate feed with clear water.
- Irrigate early in the day so foliage is dry by nightfall to prevent diseases.

STAGE 3 - Growth and development of true leaves (10-14 days)

- Soil temperature 65-70°F (18-21°C).
- Allow the soil to dry thoroughly between irrigations but avoid permanent wilting to promote root growth and control shoot growth.
- Maintain soil pH 5.5-5.8 and EC to less than 0.75 mmhos/cm.
- Increase feed to 100 150 ppm N from 20 10 20 alternating with 14 0 14 or 15-5-15 or other calcium/potassium nitrate fertilizer.
- Fertilize every 2nd 3rd irrigations.

- If using 15-0-15 supplement with magnesium 1 2x during this stage, using magnesium sulfate (16 oz/100 gal) or magnesium nitrate. Do not mix magnesium sulfate with calcium nitrate as precipitate will form!
- Use DIF whenever possible, especially the first 2 hours after sunrise, to control plant height.
- Growth regulators cannot be used on eggplant.

STAGE 4 - Plants ready for transplanting or shipping (7 days)

- Soil temperature 60-62°F (16-17°C).
- Allow soil to dry thoroughly between irrigations.
- Maintain soil pH 5.5-5.8 and EC to less than 0.75 mmhos/cm.
- Fertilize with 14 0 14 or 15-5-15 or calcium/potassium nitrate feed at 100 150 ppm N as needed.

Growing On to Finish

TEMPERATURE

- **Night:** 55-65°F (13-18°C)
- **Day:** 60-70°F (16-21°C)

LIGHT

Maintain light levels as high as possible while maintaining moderate temperatures.

MEDIA

Use a well-drained, disease-free soil-less medium with a medium initial nutrient charge and a pH 5.5-6.2.

FERTILIZATION

- Fertilize every other irrigation with 15-0-15 or 15-5-15 alternating with 20-10-20 at 150-200 ppm nitrogen.
- Maintain medium electrical conductivity around 1.0 mmhos/cm (using 1:2 extraction).

CONTROLLING HEIGHT

- Once plants are rooted to the sides of the containers they can be allowed to wilt prior to irrigation to provide some height control.
- Height can also be controlled by withholding fertilizer, especially phosphorous and ammonium-form nitrogen. Eggplant is responsive to day/night temperature differential (DIF), and is shorter with a negative DIF.

- Although Sumagic is labeled for use on Eggplant in many states, we do not recommend its use until the grower has at least trialed it on a limited basis. Application rates, timing and variety response may affect growth for the home gardener.

COMMON PROBLEMS

Insects: Aphid, Thrip, Whitefly

Diseases: Botrytis, Pythium

Post Production Care

TEMPERATURE

Night: 55-65°F (13-18°C):

Day: 60-70°F (16-21°C)

- Optimum conditions may be difficult to maintain, especially if plants are displayed outside.

LIGHT

Eggplant prefer full sun, however partial shading may be beneficial during retail display.

